VOL. 594, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY MARCH 6, 1992

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CRNS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

General

Improved equations for the calculation of chromatographic figures of merit for ideal and skewed chromatographic peaks
by M. S. Jeansonne and J. P. Foley (Baton Rouge, LA, USA) (Received July 24th, 1991) 1

Column Liquid Chromatography

Modelling of displacement chromatography using non-ideal isotherms
by P. K. de Bokx, P. C. Baarslag and H. P. Urbach (Eindhoven, Netherlands) (Received November 5th, 1991) . . . 9

High-performance liquid chromatographic study of the correlation between the physico-chemical parameters of furan and benzene aldehydes and the rate of their metabolism by yeast
by V. D. Nemirovskii, N. I. Monakhova and V. G. Kostenko (St. Petersburg, Russia) (Received October 29th, 1991) . 23

Determination of solute–polymer interaction constants via chromatography with polymer capillaries
by D. R. Jenke (Round Lake, IL, USA) (Received October 31st, 1991) 29

Retention behaviour of β-cyclodextrin complexes of anthracene and pyrene using reversed-phase liquid chromatography. The effects of *tert.*-butanol and cyclopentanol
by V. C. Anigbogu, A. Muñoz de la Peña, T. T. Ndou and I. M. Warner (Atlanta, GA, USA) (Received November 8th, 1991) 37

Selection of robust combinations of extraction liquid composition and internal standard. Monte Carlo simulation of improvement of assay methods with liquid–liquid extraction prior to high-performance liquid chromatography
by J. Wieling (Zuidlaren and Groningen, Netherlands), P. M. J. Coenegracht (Groningen, Netherlands), C. K. Mensink and J. H. G. Jonkman (Zuidlaren, Netherlands) and D. A. Doornbos (Groningen, Netherlands) (Received October 31st, 1991) 45

Computer simulation as a tool for the rapid optimization of the high-performance liquid chromatographic separation of a tryptic digest of human growth hormone
by R. C. Chloupek and W. S. Hancock (San Francisco, CA, USA) and L. R. Synder (Walnut Creek, CA, USA) (Received November 14th, 1991) 65

Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides
by B.-Y. Zhu, C. T. Mant and R. S. Hodges (Edmonton, Canada) (Received October 22nd, 1991) 75

Suppression of the effect of metal impurities in alkylsilylated silica packing materials
by K. Kimata, N. Tanaka and T. Araki (Kyoto, Japan) (Received December 9th, 1991) 87

Ligand–receptor interactions in affinity cell partitioning. Studies with transferrin covalently linked to monomethoxypoly(ethylene glycol) and rat reticulocytes
by C. Delgado, P. Sancho, J. Mendieta and J. Luque (Madrid, Spain) (Received October 29th, 1991) 97

Liquid chromatographic separation of all-carbon molecules C_{60} and C_{70} with multi-legged phenyl group bonded silica phases
by K. Jinno, K. Yamamoto, T. Ueda, H. Nagashima and K. Itoh (Toyohashi, Japan) and J. C. Fetzer and W. R. Biggs (Richmond, CA, USA) (Received October 23rd, 1991) 105

Liquid chromatographic isolation and structural elucidation of methoxymethylene dimethyl dodecatrienes catalytically synthesized from methanol and three moles of isoprene
by C. Tröltzsch and R. Berger (Greifswald, Germany) and M. Michalik (Rostock, Germany) (Received October 21st, 1991) 111

(Continued overleaf)

Contents (continued)

Identification of the constituent flavanoid units in sainfoin proanthocyanidins by reversed-phase high-performance liquid chromatography
by M. R. Koupai-Abyazani, J. McCallum and B. A. Bohm (Vancouver, Canada) (Received October 30th, 1991) . . . 117

Chromatographic separation of polyols by ligand exchange. Effects of the ion-exchange resin cross-linking and size
by H. Caruel, P. Phemius, L. Rigal and A. Gaset (Toulouse, France) (Received November 7th, 1991) 125

Determination of the norlignan glucosides of Hypoxidaceae by high-performance liquid chromatography
by P. Betto, R. Gabriele and C. Galeffi (Rome, Italy) (Received October 31st, 1991) 131

High-performance liquid chromatography of okadaic acid and free fatty acids in mussels
by F. Zonta (Trento, Italy), B. Stancher, P. Bogoni and P. Masotti (Trieste, Italy) (Received October 30th, 1991) . . 137

Isolation of aristolochic acids HA_sI and HA_sII by preparative liquid chromatography
by B. Makuch, K. Gazda and W. Cisowski (Gdańsk, Poland) (Received November 8th, 1991) 145

Determination of taxol by high-performance liquid chromatography–thermospray mass spectrometry
by S. O. K. Auriola, A.-M. Lepistö, T. Naaranlahti and S. P. Lapinjoki (Kuopio, Finland) (Received October 30th, 1991) 153

Reversed-phase liquid chromatographic behaviour of alkylamines with amperometric detection and its application to trace analysis
by M. Maruyama and T. Nagayoshi (Tokyo, Japan) (Received October 11th, 1991) 159

Chromatography of 99 amino acids and other ninhydrin-reactive compounds in the Pickering lithium gradient system
by J. A. Grunau and J. M. Swiader (Urbana, IL, USA) (Received September 10th, 1991) 165

Determination of glutamine, glutamic acid and pyroglutamic acids using high-performance liquid chromatography on dynamically modified silica
by B. Polanuer, A. Sholin, N. Demina and N. Rumiantseva (Moscow, Russia) (Received November 7th, 1991) . . . 173

High-performance gel permeation chromatographic analysis of immunoglobulin M produced by hybridoma cell culture
by N. Chauret (Quebec, Canada), J. Côté, J. Archambault and G. André (Montreal, Canada) (Received October 31st, 1991) 179

Determination of ascorbic and dehydroascorbic acid in potatoes (*Solanum tuberosum*) and strawberries using ion-exclusion chromatography
by W. D. Graham and D. Annette (Belfast, UK) (Received October 4th, 1991) 187

Solvent degradation of cloxacillin *in vitro*. Tentative identification of degradation products using thermospray liquid chromatography–mass spectrometry
by K. L. Tyczkowska (Raleigh, NC, USA), R. D. Voyksner (Research Triangle Park, NC, USA) and A. L. Aronson (Raleigh, NC, USA) (Received October 21st, 1991) 195

Determination of dirithromycin purity and related substances by high-performance liquid chromatography
by B. A. Olsen, J. D. Stafford and D. E. Reed (Lafayette, IN, USA) (Received November 4th, 1991) 203

Development of stability indicating assay methods for the determination of hydroxamic acids in topical formulations
by J. E. DiNunzio and R. R. Gadde (Buffalo, NY, USA) (Received October 22nd, 1991) 209

Determination of the mean ionic charge of the components of three ^{99m}Tc bone scanning agents by gel chromatography with two eluents of different electrolyte composition
by W. J. Gelsema, C. L. de Ligny, Y. M. Huigen and J. Schuring (Utrecht, Netherlands) (Received October 23rd, 1991) 217

Determination of major ions in snow and ice cores by ion chromatography
by C. F. Buck, P. A. Mayewski, M. J. Spencer, S. Whitlow, M. S. Twickler and D. Barrett (Durham, NH, USA) (Received October 29th, 1991) 225

Gas Chromatography

Hydrogen bonding. XXI. Solvation parameters for alkylaromatic hydrocarbons from gas–liquid chromatographic data
by M. H. Abraham and G. S. Whiting (London, UK) (Received September 4th, 1991) 229

Preparation and capillary column gas chromatographic characterization of aza-crown ether polysiloxane stationary phases
by C. Wu, J. Cheng, W. Gao, Z. Zeng, X. Lu and S. Gong (Wuhan, China) (Received October 18th, 1991) 243

Simultaneous determination of sorbic acid, dehydroacetic acid and benzoic acid by gas chromatography–mass spectrometry
by M. Kakemoto (Tokyo, Japan) (Received October 30th, 1991) 253

Analysis of zapatera olives by gas and high-performance liquid chromatography
by A. Montaño, A. de Castro, L. Rejano and A.-H. Sánchez (Seville, Spain) (Received October 29th, 1991) 259

Gas chromatographic elution patterns of chlorinated dioxins *versus* column polarity
by J. R. Donnelly and G. W. Sovocool (Las Vegas, NV, USA) (Received September 20th, 1991) 269

Determination of Pb^{2+} in water by isotope dilution gas chromatography–mass spectrometry of tetraethyllead formed by reaction with sodium tetraethylborate
by B. J. Feldman, H. Mogadeddi and J. D. Osterloh (San Francisco, CA, USA) (Received September 30th, 1991) . . 275

Supercritical Fluid Chromatography

Synthesis of (1*R-trans*)-N,N′-1,2-cyclohexylenebisbenzamideoligodimethylsiloxane copolymers for use as chiral stationary phases for capillary supercritical fluid chromatography
by D. F. Johnson, J. S. Bradshaw, M. Eguchi, B. E. Rossiter and M. L. Lee (Provo, UT, USA) and P. Petersson and K. E. Markides (Uppsala, Sweden) (Received November 18th, 1991) 283

Correlation between column surface area and retention of polar solutes in packed-column supercritical fluid chromatography
by T. A. Berger and J. F. Deye (Avondale, PA, USA) (Received October 30th, 1991) 291

Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies
by J. J. Langenfeld, M. D. Burford, S. B. Hawthorne and D. J. Miller (Grand Forks, ND, USA) (Received October 18th, 1991) 297

Supercritical carbon dioxide extraction of resin and fatty acids from sediments at pulp mill sites
by H.-B. Lee and T. E. Peart (Burlington, Canada) (Received November 5th, 1991) 309

Electrophoresis

Charge-reversed, polymer-coated capillary column for the analysis of a recombinant chimeric glycoprotein
by K. Tsuji and R. J. Little (Kalamazoo, MI, USA) (Received November 12th, 1991) 317

Capillary modification and evaluation using streaming potential and frontal chromatography for protein analysis in capillary electrophoresis
by T. Wang and R. A. Hartwick (Binghamton, NY, USA) (Received November 1st, 1991) 325

Separation of plant growth regulators by capillary electrophoresis
by S. K. Yeo, H. K. Lee and S. F. Y. Li (Kent Ridge, Singapore) (Received October 18th, 1991) 335

Finite difference modelling of continuous-flow electrophoresis
by T. M. Grateful and E. N. Lightfoot, Jr. (Madison, WI, USA) (Received November 12th, 1991) 341

Continuous-flow electrophoresis: a separation criterion applied to the separation of model proteins
by H. Roux-de Balmann and V. Sanchez (Toulouse, France) (Received October 8th, 1991) 351

Continuous separation and purification of proteins by recycling isotachophoresis
by J. Caslavska and W. Thormann (Berne, Switzerland) (Received August 27th, 1991) 361

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. III. Analysis of a crude rare earth chloride from monazite
by J. Hu, T. Hirokawa, F. Nishiyama, Y. Kiso, K. Ito and E. Shoto (Higashi-Hiroshima, Japan) (Received October 31st, 1991) 371

Planar Chromatography

Quantitative separation of Cr^{3+} from Mo^{6+}, W^{6+}, Hg_2^{2+}, Cu^{2+} and Pb^{2+}. Chromatographic behaviour of 51 cations on papers impregnated with Sn(IV)-based inorganic ion exchangers in complex-forming acid systems
by S. D. Sharma and S. Misra (Moradabad, India) (Received November 6th, 1991) 379

(Continued overleaf)

Contents (continued)

SHORT COMMUNICATIONS

Column Liquid Chromatography

Primary and secondary amine derivatization with luminarins 1 and 2: separation by liquid chromatography with peroxyoxalate chemiluminescence detection
by M. Tod (Bobigny, France), J.-Y. Legendre (Paris, France), J. Chalom (Les Ulis, France) and H. Kouwatli, M. Poulou, R. Farinotti and G. Mahuzier (Châtenay-Malabry, France) (Received November 21st, 1991) 386

Fast protein monitoring in fermentation both using non-porous micropellicular reversed-phase columns
by E. Watson and F. Yao (Thousand Oaks, CA, USA) (Received November 18th, 1991) 392

Gas Chromatography

Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection
by A. Sarafraz-Yazdi, M. Y. Khuhawar and P. C. Uden (Amherst, MA, USA) (Received December 10th, 1991) . . . 395

Electrophoresis

Fast and sensitive staining technique for catalase in polyacrylamide gel
by H. Pichorner, S. Loew, S. A. A. Korori, G. Jessner and R. Ebermann (Vienna, Austria) (Received November 5th, 1991) 400

BOOK REVIEW

Advances in steroid analysis '90 (edited by S. Görög; guest editor E. Heftmann), reviewed by T. Nambara (Sendai, Japan) . 403

Author Index 404

Erratum 407

Announcement of Special Issue on Principles and Applications of Solid Phase Extraction 408

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*).